



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**16.10.2002 Bulletin 2002/42**

(51) Int Cl.7: **A23J 3/08**, A23L 1/305,  
A23C 19/09, A23C 19/082,  
A23L 1/314, A23D 7/00

(21) Application number: **01107852.4**

(22) Date of filing: **09.04.2001**

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE TR**  
Designated Extension States:  
**AL LT LV MK RO SI**

• **Spiegel, Thomas L., Dr.-Ing.**  
**81737 München (DE)**

(71) Applicant: **Kraft Foods R&D, Inc.**  
**81737 Munich (DE)**

(74) Representative: **HOFFMANN - EITLE**  
**Patent- und Rechtsanwälte**  
**Arabellastrasse 4**  
**81925 München (DE)**

(72) Inventors:  
• **Wolfschoon Pombo, Alan F., Dr.-Ing.**  
**81737 München (DE)**


Remarks:

Amended claims in accordance with Rule 86 (2)  
EPC.

(54) **Process for incorporating whey proteins into foodstuffs**

(57) The present invention is concerned with the incorporation of whey proteins into foodstuffs by acidifying an aqueous solution of one or more whey proteins below their isoelectric pH, optionally forming a whey protein-stabilized fatty emulsion by blending and homogenizing said acidified solution of whey proteins with one or more

fats, heat-treating said acidified solution or whey protein-stabilized fatty emulsion and blending same with a foodstuff base to form a foodstuff, and optionally blending and incubating the foodstuff with a transglutaminase.



# Description

[0001] The present invention is concerned with the incorporation of whey proteins into foodstuffs. In particular, the invention relates to a process for preparing a whey protein-stabilized fatty emulsion, the thus obtainable whey protein-stabilized fatty emulsion, a process for incorporating whey proteins into a foodstuff using said emulsion or an acidified solution of whey proteins, and the foodstuff thus obtainable.

[0002] In the field of the manufacture of food, several attempts have been made to introduce larger amounts of whey proteins, for example to replace costly casein proteins and utilize wasted but rich whey proteins, without a resulting gumminess (texture effect). With respect to the example of process cheese, it has been practically impossible to incorporate larger amounts of whey proteins over a weight ratio of whey protein to casein of about 10:90. The higher the amount of whey proteins, the gummier and softer is the resultant cheese.

[0003] The following documents, the disclosure of which is incorporated herein by reference, represent technological background material on whey proteins, their properties and usage in foods:

Kinsella, J.E.; Whitehead, D.M. (1989): Proteins in whey: Chemical, physical, and structural properties. Advances in Food and Nutrition Research, 33, 343-438

Konrad, G.; Lieske, B. (1994): Gezielte thermische Denaturierung - ein alternativer Weg zu funktionellerem Molkenprotein. Deutsche Milchwirtschaft, 45, 1130-1134

Lupano, C. (1994): Effect of heat treatments in very acidic conditions on whey protein isolate properties. Journal of Dairy Science, 77, 2191-2198

Patocka, G.; Drathen, M.; Jelen, P. (1987): Heat stability of isolated whey protein fractions in highly acidic conditions. Milchwissenschaft, 42, 700-705

Shimizu, M.; Saito, M.; Yamauchi, K. (1985): Emulsifying and structural properties of  $\beta$ -lactoglobulin at different pHs. Agricultural and Biological Chemistry, 49, 189-194

Strandholm, J.J.; Prochnow, R.R.; Miller, M.S.; Woodford, L.E.; Naunaber, S.M. (1989): Method for controlling melting properties of process cheese. Patent US 4 885 183

Swartz, M.L. (1988): Beverage cloud based on a whey protein-stabilized lipid. Patent US 4 790 998

Yamauchi, K.; Shimizu, M.; Kamiya, T. (1980): Emulsifying properties of whey protein. Journal of Food Science, 45, 1237-1242

[0004] As regards the structural and textural properties of foods, one possibility to increase the firmness thereof would be the polymerization of the proteins by an enzyme, namely a transglutaminase. However, transglutaminases predominantly polymerize caseins, whereas their reactivity towards whey proteins is very low or even absent. Thus, the effect of enzymatic treatment on whey protein-enriched food compositions is considerably low.

[0005] The following documents represent technological background material on transglutaminases, their effect on proteins and usage in foods:

Aboumahmoud, R.; Savello, P. (1990): Cross-linking of whey proteins by transglutaminase. Journal of Dairy Science 73, 256-263

Budolfsen, G.; Nielsen, P.M. (1999): Method for production of an acidified edible gel on milk basis. Patent US 5 866 180

Chanyongvorakul, Y.; Matsumara, Y.; Sawa, A.; Nio, N.; Mori, T. (1997): Polymerization of  $\beta$ -lactoglobulin and bovine serum albumin at oil-water interfaces in emulsions by transglutaminase. Food Hydrocolloids 11, 449-455

Faergemand, M.; Murray, B.S.; Dickinson, E. (1997): Cross-linking of milk proteins at the oil-water interface. Journal of Agricultural and Food Chemistry 45, 2514-2519

Faergemand, M.; Otte, J.; Qvist, K.B. (1997): Enzymatic cross-linking of whey proteins by a  $\text{Ca}^{2+}$ -independent microbial transglutaminase from *Streptomyces lydicus*. Food Hydrocolloids 11, 19-25

Feargemand, M.; Otte, J.; Qvist, K.B. (1997): Emulsifying properties of milk proteins cross-linked with microbial transglutaminase. *International Dairy Journal* 8, 715-723

Kuraishi, T.; Sakamoto, J.; Soeda, T. (1996): Production of cheese using transglutaminase. Patent Application EP 0 711 504 (JP 81-73032 A2)

Soeda, T. (1999): Cheese whey protein having improved texture, process for producing the same and use thereof. Patent Application EP 0 966 887

Traoré, F.; Meunier, J.C. (1992): Cross-linking activity of placental FXIIIa on whey proteins and caseins. *Journal of Agricultural and Food Chemistry* 40, 399-402

Tsukasaki, F.; Minagawa, E.; Mikami, T.; Nonaka, M.; Motoki, M. (1990): Preparation of cheese food. Patent JP 21-31537 A2

Yamamoto, Y. (1996): Rheology of milk protein gels and protein-stabilized emulsions cross-linked by transglutaminase. *Journal of Agricultural and Food Chemistry* 44, 1371-1377

[0006] In view of the above, the present inventors thoroughly studied the properties of aqueous solutions of whey proteins and conceived new ways to modify such proteins, thus providing a solution to the problems of the prior art as outlined above.

[0007] Accordingly, in a first aspect, the present invention provides a process for preparing a whey protein-stabilized fatty emulsion, comprising the steps of

- acidifying an aqueous solution of one or more whey proteins below their isoelectric pH,
- blending and homogenizing the acidified solution of whey proteins with one or more fats to form a whey protein-stabilized fatty emulsion, and
- heat-treating said whey protein-stabilized fatty emulsion at a temperature of more than 80 °C.

[0008] Potential sources for the whey proteins referred to herein are whey protein concentrates (WPC) and whey protein isolates (WPI), either as reconstituted WPC and/or WPI powders or preferably as liquid concentrates such as ultrafiltrated whey. Likewise, normal whey powder may represent a source for the whey proteins referred to in this description. The whey can be rennet whey or acid whey. Microfiltration permeate obtained by using, e.g., a 0.1 µm membrane, which permeate contains the native whey proteins, and the ultrafiltrated concentrate thereof may also be used. A combination of one or more whey protein sources may be utilized to provide whey proteins in the present invention.

[0009] The whey proteins initially used in the present invention may be native or denatured whey proteins, with preference given to predominantly native whey proteins. However, particulate and highly denatured whey proteins also exhibit very good effects in the present invention.

[0010] The protein content in the aqueous solution of one or more whey proteins is preferably at least 4 % by weight (with total solids being, for example, about 10 % or more) to reach the desired whey protein content in the final food products. The maximum whey protein content is preferably about 20 % by weight. More preferably, the whey protein content of the aqueous solution is in the range of 8 to 12 % by weight (preferably 12 to 20 % total solids).

[0011] In the first step of the process for preparing a whey protein-stabilized fatty emulsion, the aqueous solution of one or more whey proteins as outlined above is acidified below the isoelectric pH of the whey proteins contained in said solution. For the purpose of acidifying the whey protein solution, any acid may be used that is not objectionable to the intended application and use of the final whey protein-stabilized fatty emulsion. Specifically, any food grade acid such as lactic, citric, phosphoric or hydrochloric acid and any acidulant such as glucono-δ-lactone or vinegar may be used, alone or in combination of two or more thereof. The preferred acid is 90 % lactic acid. Optionally, it is possible to employ a lactic acid producing bacterial culture in the presence of a suitable sugar such as glucose or lactose, optionally in combination with one or more of the acids mentioned before. In general, the concentration of the acid(s) and the temperature used during acidification are adjusted such that no detrimental effects are induced. For example, if there are caseins present in the solution, it may be necessary to adjust the initial concentration of the acid(s) and the temperature during acidification in such a way that no coagulation (flocculation) of caseins is induced. Similarly, too high (e.g. >65 °C) temperatures may induce a whey protein coagulation if the whey proteins are in the native state or not fully denatured. Accordingly, room (ambient) temperature represents a preferred temperature for a practical application.

[0012] The pH accomplished in the acidification step is preferably considerably below the isoelectric point of the

whey proteins comprised in the aqueous solution. Thus, the pH reached is preferably in the range of 4.5 to 2.5, more preferably 4.0 to 3.5. As a reference, the isoelectric points of major whey proteins are: ~5.1 ( $\beta$ -lactoglobulin); 4.2-4.5 ( $\alpha$ -lactalbumin); 4.7-4.9 (bovine serum albumin) and 6.3-7.0 (immunoglobulin G1).

**[0013]** In the acidification step, the aqueous solution of the whey proteins is preferably agitated when adding the acid in order to avoid local peak concentrations of the acid. Although it is also possible to add the aqueous solution of whey proteins to an aqueous solution of the acid(s), this may not be the preferred way of acidification due to potential detrimental peak acid concentrations. Preferably, the whey protein solution after addition of the acid is allowed to stand or is agitated until an equilibrium state is reached, for example, for one or more minutes, preferable about 10 to 30 minutes.

**[0014]** In the second step of the process for preparing a whey protein-stabilized fatty emulsion, the acidified solution of whey proteins is blended and homogenized with one or more fats. In this step, any fat may be used which is not objectionable to the desired application of the resulting fatty emulsion. The preferred fat is milk fat, e.g. in the form of cream, plastic cream and more preferably butter or anhydrous butter fat. However, other sources for fats may be used, especially vegetable fats and oils, or animal fat, such as beef tallow, depending on the application of the whey protein-stabilized fatty emulsion. A combination of two or more fats may be used. The one or more fats are used in an amount such that the ratio of whey protein to fat in the resulting fatty emulsion is in the range of preferably 3:1 to 1:5 (the total fat content in the fatty emulsion being preferably in the range of 3 to 30 % by weight), more preferably 2:1 to 1:2.

**[0015]** The blending of the acidified solution of whey proteins with one or more fats is followed by a homogenization, i.e. a shear treatment. Blending and homogenizing the two components, i.e. acidified whey protein solution and fats, may be a combination of one or more steps. For example, the components may first be blended and subsequently subjected to a shear treatment, or blending may be effected simultaneously in the homogenization, for example by introducing the components into a homogenizer and starting the device. Preferably, blending and homogenizing is carried out in a single apparatus for economic reasons and effectiveness. The blending and homogenizing step may be carried out using a conventional homogenizer and is preferably carried out at a temperature of from 50 to 70 °C and a pressure of from 100 to 300 bar, depending on the fat content (lower pressures may be more suitable at higher fat levels). In general, the homogenization can be conducted at a temperature and pressure and for a period which are conventional in this art.

**[0016]** It is assumed that the resulting whey protein-stabilized fatty emulsion is of the oil-in-water emulsion type. However, this assumption should not be construed as restricting the scope of the present invention, and other structures such as water-in-oil or bicontinuous structures may be contemplated, depending on the components and amounts used thereof.

**[0017]** The blending and homogenizing step is followed by a heat treatment at a temperature of 80 °C or more, and optionally at an elevated pressure. The preferred temperature range for the heat treatment is 80 to 95 °C, and the preferred holding time is in the range of 1 to 10 minutes, for example a heat treatment condition of a temperature of 85 °C and a holding time of 5 minutes. It is assumed that the subsequent heat treatment should modify the topography of the already changed whey proteins (due to the acidification below their isoelectric pH) to make them even more functional, for example in replacing casein in casein-containing foodstuffs.

**[0018]** Following the blending/homogenization or the heat treatment, the whey protein-stabilized fatty emulsion may be cooled (e.g., to room temperature) or directly used in the hot state in the desired application such as outlined below.

**[0019]** The whey protein-stabilized fatty emulsion obtainable by the process of the present invention preferably has a fat content in the range of from 3 to 30 % by weight and a whey protein content in the range of from 5 to 15 % by weight. In a preferred embodiment, an emulsion made in accordance with the process of the invention (e.g., by mixing WPC and butter) has 20 to 25 % total solids, 5 to 10 % fat, 8 to 12 % protein and 3 to 5 % sugars, based on the total weight of the emulsion (the presence of additional components such as sugars may result from the whey protein source such as WPC or the deliberate addition of further components).

**[0020]** The whey protein-stabilized fatty emulsion obtainable by the above process represents another aspect of the present invention.

**[0021]** The protein-stabilized fatty emulsion of the invention may be used for incorporating whey proteins into a foodstuff. Thus, in another aspect the present invention provides a process for incorporating whey protein into a foodstuff comprising the steps of blending the whey protein-stabilized fatty emulsion obtainable by the above process of the invention with a foodstuff base to form a foodstuff. In an alternative embodiment, the process for incorporating whey proteins into a foodstuff comprises the steps of acidifying an aqueous solution of one or more whey proteins below their isoelectric pH, heat-treating the acidified solution at a temperature of more than 80 °C and blending the acidified and heat-treated solution with a foodstuff base to form a foodstuff. In this alternative embodiment, the aspects, conditions and properties of acidifying an aqueous solution of one or more whey proteins below their isoelectric pH and heat-treating the acidified solution are as defined above having regard to the process for preparing a whey protein-stabilized fatty emulsion.

**[0022]** The foodstuff base may be any dairy or non-dairy based foodstuff base. In fact, the present inventors have

found that using a specifically acidified and heat-treated whey protein solution or whey protein fatty emulsion provides a means for using higher amounts of whey proteins, or to incorporate more whey proteins, into other food systems. Preferably, the foodstuff base is a dairy based foodstuff base and more preferably a casein-containing foodstuff base. Moreover, the foodstuff base can be foodstuff base containing meat proteins. Specific examples of the foodstuff base are bases for cream cheese, process cheese, natural cheese and mayonnaise, as well as process meat products.

**[0023]** The conditions for blending the fatty emulsion and/or the acidified and heat-treated whey protein solution and the foodstuff base are those normally found in the manufacture (blending of raw materials) of foodstuffs, for example process cheese, cream cheese or natural cheese, or process meat. If the whey protein-stabilized fatty emulsion of the invention is used, the weight ratio of fatty emulsion to foodstuff base is preferably in the range of from 20:80 to 70:30. Similarly, if the acidified and heat-treated solution of one or more whey proteins is used, the weight ratio of whey proteins to foodstuff base is preferably in the range of from 20:80 to 70:30. If a casein-containing foodstuff base is used, the blending ratio of fatty emulsion or acidified whey protein solution to casein-containing foodstuff base is preferably such that the weight ratio of whey proteins to casein in the resulting casein-containing foodstuff is in the range of >10:90 to 80:20, more preferably 20:80 to 40:60. If a meat protein-containing foodstuff base is used, the blending ratio of fatty emulsion or acidified whey protein solution to meat protein-containing foodstuff base is preferably such that the weight ratio of whey proteins to meat proteins in the resulting meat protein-containing foodstuff is in the range of from 10:90 to 80:20, more preferably 20:80 to 40:60. Especially, if meat proteins are used, the resulting meat protein-containing foodstuff can also include animal fat, such as beef tallow.

**[0024]** If desired, the resulting foodstuff may be further blended and incubated with a transglutaminase enzyme. Normally, whey proteins do not react with transglutaminases. The present inventors have found that the pre-treatment of whey protein solutions in the acid pH range (below the isoelectric pH of the whey proteins) as described in this invention offers a possibility to modify the behavior of whey proteins also towards the action of transglutaminase. Accordingly, in another aspect of the present invention, the process for incorporating whey proteins into a foodstuff comprises the further step of blending and incubating the resulting foodstuff with the enzyme transglutaminase. As is generally known by the person skilled in the art, there are different transglutaminases, classified under the general nomenclature EC 2.3.2.13, and any of these transglutaminases can be used in the present invention. In particular, it is possible to use a transglutaminase produced by *Streptovorticillium mobaraense* which is commercially available under the name of "Aktiva-MP" from Ajinomoto. The enzyme employed is preferably used in conventional amounts such as 1 to 6 u/g (units enzyme per gram protein in the system). The incubating conditions are preferably a temperature in the range of 20 to 60 °C, more preferably 50 °C; a duration of 5 to 60 minutes; and a pH of 5.5 to 7.5, more preferably 6 to 7. Preferably, no shear is applied during the incubation period.

**[0025]** The foodstuff which is obtainable by the process according to the invention, for example process cheese, cream cheese, natural cheese, mayonnaise or process meat, is superior as compared to foodstuffs of the prior art in that a high firmness and short texture of the resultant products can be accomplished even at high whey protein contents, which is not observed when native whey proteins are utilized. Specifically, if a casein-containing foodstuff is manufactured, such as a process cheese formulation, high protein to casein ratios of up to 80:20 by weight can be established together with a favorable appearance and excellent sensory properties.

#### Example 1

**[0026]** A whey protein concentrate (17.7 % total solids, 10 % protein) with the majority of its proteins in their native state is acidified with lactic acid down to pH 3.8. Afterwards the acidified whey protein solution is blended with 5 % molten butter at room temperature and homogenized at 220 bar and 60 °C. The emulsion is then heat treated at 85 °C for 5 min. Then 34.2 parts of the heated emulsion are immediately blended with 3.5 parts of rennet casein, 3.5 parts of milk concentrate, 1.4 parts of starch, 7 parts of hard cheese, 15.9 parts of butter, 5.3 parts of whey powder, 3.0 parts of emulsifying salts, 1 part of salt and 25.2 parts of water so as to complete 100 parts. After thoroughly mixing, the blend is treated as in the conventional manufacture of process cheese spreads.

#### Example 2

**[0027]** A whey protein concentrate (17.7 % total solids, 10 % protein) with the majority of its proteins in their native state is acidified with lactic acid down to pH 3.8. Afterwards the acidified whey protein solution is blended with 5 % molten butter at room temperature and homogenized at 220 bar and 60 °C. The emulsion is then heat treated at 85 °C for 5 min. Then 34.2 parts of the heated emulsion are immediately blended with 3.5 parts of rennet casein, 3.5 parts of milk concentrate, 1.4 parts of starch, 7 parts of hard cheese, 15.9 parts of butter, 5.3 parts of whey powder, 3.0 parts of emulsifying salts, 1 part of salt and 25.2 parts of water so as to complete 100 parts. After thoroughly mixing, the blend is incubated with a microbial transglutaminase (5 units per gram protein in the blend) at 50 °C for 1 h. The blend is then treated as in the conventional manufacture for process cheese spreads. The following table shows the firmness

(Stevens Texture Analyzer) of the final cheese produced as described above compared to a standard product and a product with a native whey protein concentrate.

Product	Whey protein: casein ratio	Transglutaminase	Firmness [g]
Standard	10:90	no	57
		yes	160
With native whey protein	45:55	no	36
		yes	64
With whey protein emulsion pH 3.8, heated	45:55	no	42
		yes	115

## Claims

1. A process for preparing a whey protein-stabilized fatty emulsion comprising  
acidifying an aqueous solution of one or more whey proteins below their isoelectric pH,  
blending and homogenizing the acidified solution of whey proteins with one or more fats to form a whey  
protein-stabilized fatty emulsion, and  
heat-treating said whey protein-stabilized fatty emulsion at a temperature of more than 80°C.
2. The process according to claim 1 wherein the weight ratio of whey proteins to fats is in the range of 3:1 to 1:5  
based on dry matter.
3. A whey protein-stabilized fatty emulsion obtainable by the process according to claim 1 or 2.
4. A process for incorporating whey proteins into a foodstuff comprising  
blending the whey protein-stabilized fatty emulsion obtainable by the process according to claim 1 or 2 with  
a foodstuff base to form a foodstuff.
5. A process for incorporating whey proteins into a foodstuff comprising  
acidifying an aqueous solution of one or more whey proteins below their isoelectric pH,  
heat-treating said acidified solution at a temperature of more than 80 °C, and  
blending the acidified solution of one or more whey proteins with a foodstuff base to form a foodstuff.
6. The process of claim 4 or 5 wherein the resulting foodstuff is further blended and incubated with a transglutaminase.
7. The process of any one of claims 4 to 6 wherein the foodstuff base contains casein.
8. The process of claim 7 wherein the weight ratio of whey proteins to casein in the resulting casein-containing food-  
stuff is in the range of >10:90 to 80:20, preferably 20:80 to 40:60.
9. The process of any one of claims 4 to 8 wherein the foodstuff base is a process cheese formulation.
10. The process of any one of claims 4 to 6 wherein the foodstuff base contains meat proteins.
11. The process of claim 10 wherein the weight ratio of whey proteins to meat proteins in the resulting meat protein-  
containing foodstuff is in the range of from 10:90 to 80:20, preferably 20:80 to 40:60.
12. The process of claim 10 or 11 wherein the resulting meat protein-containing foodstuff includes animal fat.
13. The process of any of claims 10 to 12 wherein the foodstuff base is process meat base.
14. A foodstuff obtainable by the process according to any one of claims 4 to 13.
15. The foodstuff according to claim 14 which is a process cheese.

16. The foodstuff according to claim 14 which is a process meat product.

**Amended claims in accordance with Rule 86(2) EPC.**

5

1. A process for preparing a whey protein-stabilized fatty emulsion comprising  
acidifying an aqueous solution of one or more whey proteins below their isoelectric pH,  
blending and homogenizing the acidified solution of whey proteins with one or more fats to form a whey  
protein-stabilized fatty emulsion, and  
10 heat-treating said whey protein-stabilized fatty emulsion at a temperature of more than 80°C.

2. The process according to claim 1 wherein the weight ratio of whey proteins to fats is in the range of 3:1 to 1:5  
based on dry matter.

15 3. A whey protein-stabilized fatty emulsion obtainable by the process according to claim 1 or 2.

4. A process for incorporating whey proteins into a foodstuff comprising  
blending the whey protein-stabilized fatty emulsion obtainable by the process according to claim 1 or 2 with  
a foodstuff base to form a foodstuff.

20

5. A process for incorporating whey proteins into a foodstuff comprising  
acidifying an aqueous solution of one or more whey proteins below their isoelectric pH,  
heat-treating said acidified solution at a temperature of more than 80 °C,  
blending the acidified solution of one or more whey proteins with a foodstuff base to form a foodstuff, and  
25 further blending and incubating the resulting foodstuff with a transglutaminase.

25

6. The process of claim 4 wherein the resulting foodstuff is further blended and incubated with a transglutaminase.

7. The process of any one of claims 4 to 6 wherein the foodstuff base contains casein.

30

8. The process of claim 7 wherein the weight ratio of whey proteins to casein in the resulting casein-containing  
foodstuff is in the range of >10:90 to 80:20, preferably 20:80 to 40:60.

9. The process of any one of claims 4 to 8 wherein the foodstuff base is a process cheese formulation.

35

10. The process of any one of claims 4 to 6 wherein the foodstuff base contains meat proteins.

11. The process of claim 10 wherein the weight ratio of whey proteins to meat proteins in the resulting meat protein-  
containing foodstuff is in the range of from 10:90 to 80:20, preferably 20:80 to 40:60.

40

12. The process of claim 10 or 11 wherein the resulting meat protein-containing foodstuff includes animal fat.

13. The process of any of claims 10 to 12 wherein the foodstuff base is process meat base.

14. A foodstuff obtainable by the process according to any one of claims 4 to 13.

45

15. The foodstuff according to claim 14 which is a process cheese.

16. The foodstuff according to claim 14 which is a process meat product.

50

55



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 01 10 7852

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InLCI.7)
X	US 4 188 411 A (KUIPERS ARIE ET AL) 12 February 1980 (1980-02-12)	5,7,9, 14,15	A23J3/08 A23L1/305 A23C19/09 A23C19/082 A23L1/314 A23D7/00
Y	* claims; examples *	6	
D,Y	EP 0 966 887 A (AJINOMOTO KK) 29 December 1999 (1999-12-29) * paragraphs '0039!', '0040!', '0059!'; claims *	6	
X	US 3 956 520 A (AIELLO JOSEPH A) 11 May 1976 (1976-05-11) * column 7 - column 8; claims *	5,7,9, 14,15	
X	US 3 930 039 A (KUIPERS ARIE) 30 December 1975 (1975-12-30) * claims; examples *	5,14	
X	US 3 922 375 A (DALAN ERNESTO ET AL) 25 November 1975 (1975-11-25) * column 3, line 19 - line 46; example 2 *	5,14	
X	WO 98 36647 A (KENNEDY BRENDAN T O ;KELLY PHILIP M (IE); TEAGASC AGRIC FOOD DEV A) 27 August 1998 (1998-08-27) * claims *	5,14	TECHNICAL FIELDS SEARCHED (InLCI.7)  A23J A23L A23C A23D
X	GB 1 440 182 A (UNILEVER LTD) 23 June 1976 (1976-06-23) * the whole document *	1-5,14	
X	EP 0 603 981 A (VER COOP MELKIND) 29 June 1994 (1994-06-29) * claims *	3,4,14 1	
X	EP 0 398 408 A (UNILEVER NV ;UNILEVER PLC (GB)) 22 November 1990 (1990-11-22) * claims; examples *	3,4,7 1	
-/--			
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>3 October 2001</b>	Examiner <b>Boddaert, P</b>
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPC FORM 1533 (3.8.92) (P-42/01)

11/020742  
11/244020





European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 01 10 7852

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	EP 0 818 149 A (SOREMARTEC SA ; FERRERO OHG (DE); FERRERO SPA (IT)) 14 January 1998 (1998-01-14)	3,4,7-9, 14,15	
A	* claims; examples *	1	
X	EP 0 787 436 A (SNOW BRAND MILK PROD CO LTD ; SNOW BRAND FOOD CO LTD (JP)) 6 August 1997 (1997-08-06)	1,3,4, 10-14,16	
A	* the whole document *		
X	US 5 350 590 A (MCCARTHY ANTHONY J ET AL) 27 September 1994 (1994-09-27)	3-5,7,9, 14,15	
A	* the whole document *	1	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 3 October 2001	Examiner Boddaert, P
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background I : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EP-0 FORM 1503 03 82 (PC/2001)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 10 7852

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

03-10-2001

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4188411	A	12-02-1980	DE 2503840 A1	05-08-1976
			BE 838032 A1	14-05-1976
			FR 2298955 A1	27-08-1976
			NL 7600815 A ,B,	03-08-1976
			SE 428519 B	11-07-1983
			SE 7600779 A	02-08-1976
EP 0966887	A	29-12-1999	JP 2000004786 A	11-01-2000
			AU 3394599 A	13-01-2000
			BR 9902614 A	16-05-2000
			EP 0966887 A1	29-12-1999
US 3956520	A	11-05-1976	NONE	
US 3930039	A	30-12-1975	DE 2138277 A1	08-02-1973
			AU 451606 B	15-08-1974
			AU 3352671 A	22-03-1973
			FR 2149039 A5	23-03-1973
			GB 1313085 A	11-04-1973
			JP 56001053 B	10-01-1981
			NL 7113312 A ,B,	01-02-1973
			NL 7215050 A	11-05-1973
			SE 384314 B	03-05-1976
US 3922375	A	25-11-1975	CH 556143 A	29-11-1974
			AU 467778 B	11-12-1975
			AU 5775173 A	09-01-1975
			CA 1011985 A1	14-06-1977
			DE 2345798 A1	21-03-1974
			FR 2198704 A1	05-04-1974
			GB 1402977 A	13-08-1975
			JP 1224945 C	31-08-1984
			JP 49062670 A	18-06-1974
			JP 58020577 B	23-04-1983
			NL 7311095 A	13-03-1974
			ZA 7304649 A	26-06-1974
WO 9836647	A	27-08-1998	IE 970105 A2	15-07-1998
			AU 6112298 A	09-09-1998
			EP 0967884 A1	05-01-2000
			WO 9836647 A1	27-08-1998
GB 1440182	A	23-06-1976	BE 506697 A	
			DE 2325133 A1	13-12-1973
			FI 55287 B	30-03-1979
			FR 2185018 A1	28-12-1973

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 01 10 7852

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

03-10-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1440182 A		FR 1048667 A	23-12-1953
		IT 500010 A	
		NL 7306716 A ,B	20-11-1973
		NL 7314932 A ,B	02-05-1974
		SE 390868 B	31-01-1977
		US 3944680 A	16-03-1976
EP 0603981 A	29-06-1994	NL 9202245 A	18-07-1994
		DE 69305312 D1	14-11-1996
		DE 69305312 T2	28-05-1997
		DK 603981 T3	17-03-1997
		EP 0603981 A1	29-06-1994
		FI 935761 A	24-06-1994
EP 0398408 A	22-11-1990	AU 5490390 A	15-11-1990
		CA 2016711 A1	15-11-1990
		EP 0398408 A2	22-11-1990
		JP 3007538 A	14-01-1991
		ZA 9003700 A	29-01-1992
EP 0818149 A	14-01-1998	CH 690351 A5	15-08-2000
		BR 9702573 A	29-09-1998
		CA 2209585 A1	08-01-1998
		CZ 9702134 A3	14-01-1998
		EP 0818149 A2	14-01-1998
		HU 9701154 A2	30-03-1998
		PL 320949 A1	19-01-1998
		SK 90097 A3	14-01-1998
EP 0787436 A	06-08-1997	EP 0787436 A1	06-08-1997
		JP 3050606 B2	12-06-2000
		US 5948462 A	07-09-1999
		WO 9704669 A1	13-02-1997
US 5350590 A	27-09-1994	NONE	

EPOFORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82